



Potentially Preventable Hospitalizations Program Surveillance Report – Trinity County June 2016

Contents

Introduction.....	2
The Texas Potentially Preventable Hospitalizations Program	3
Definitions	4
Key Findings.....	5
Demographics (Race/Ethnicity and Gender), Trinity County and Potentially Preventable Hospitalizations (PPH), 2014	6
Table 1 – Trinity County Demographics by Race/Ethnicity and Gender, 2014	6
Table 2 – PPH Demographics by Race/Ethnicity and Gender, 2014.....	6
Figure 1 - Percentage of Trinity County Adults by Race/Ethnicity, 2014	7
Figure 2 - Percentage of PPH by Race/Ethnicity, Trinity County, 2014	7
Demographics (Age Groups), Trinity County and Potentially Preventable Hospitalizations (PPH), 2014.....	8
Table 3 – Trinity County Demographics by Age Groups, 2014	8
Table 4 – PPH Demographics by Age Groups, Trinity County, 2014	8
Number of Potentially Preventable Hospitalizations (PPH) and Risk Adjusted Rates, 2014.....	9
Table 5 - Number of PPH and the Risk Adjusted Rates, Trinity County, 2014.....	9
Figure 3 – Risk Adjusted Rates of PPH, Trinity, 2014.....	10
Figure 4 – Risk Adjusted Rates of PPH for 4 Selected Conditions, Trinity County, 2014.....	11
Hospital Charges and Length of Stay of Potentially Preventable Hospitalizations (PPH), Trinity County, 2014.....	15
Table 6 – Hospital Charges and Length of Stay of PPH, Trinity County, 2014.....	15
Expected Primary Source of Payment of Potentially Preventable Hospitalizations (PPH), Trinity County, 2014 ...	16
Table 7 – Expected Primary Payer of PPH, Trinity County, 2014	16
Future Recommendations	17
Technical Notes	17

Measure Information:	17
Limitations:	17
Related Links:.....	18

Introduction

Potentially Preventable Hospitalizations (PPH) are hospital admissions for certain acute illnesses and chronic conditions that may be avoided with appropriate outpatient treatment and disease management. Lack of access to healthcare and poor-quality care lead to increases in these types of hospitalizations. PPHs are also referred to as *Ambulatory Care Sensitive Conditions*, *Prevention Quality Indicators* (PQIs), and *Potentially Preventable Admissions/Events*.

Methodology to identify PPHs was developed by the Agency for Healthcare Research and Quality (AHRQ). AHRQ is the lead federal agency responsible for research on healthcare quality costs, outcomes and patient safety. The hospitalizations are geographically identified by the residence of the patient—not the location where they were hospitalized.

PPH illnesses and conditions are identified by their primary diagnosis in the hospital. The most common chronic conditions that result in preventable hospitalizations are: angina, congestive heart failure (CHF), hypertension, chronic obstructive pulmonary disease (COPD) and asthma in older adults, diabetes short-term complications, and diabetes long-term complications. The most common acute illnesses that result in preventable hospitalizations are: bacterial pneumonia, dehydration, and urinary tract infection (UTI).

This report analyzes **adult** potentially preventable hospitalizations data. Hospitalizations of children are not included. Not all of the PPH conditions identified by AHRQ are included in this report. For a complete list, please visit the AHRQ website. (See page 18) PPH chart and table totals appearing in this report are not meant to be utilized as composite indicators.

PPH data and trends may be used as tools to improve the outpatient healthcare system and population health. Although these indicators are based on hospital inpatient data, they provide insight into community resources and services outside the hospital setting, and are markers of health system efficiency and efficacy. This information is not meant to be used as an evaluation of particular hospitals or other healthcare providers.

The Texas Potentially Preventable Hospitalizations Program

DSHS approved a Project Charter in January 2008 to support community assessment projects using PPH data. An Exceptional Item proposal was developed in 2010 to provide resources to reduce PPH rates. In June of 2011, the 82nd Texas Legislature approved \$2 million for preventative hospitalization projects through a rider to the appropriations bill.

In August of 2011, DSHS sent a Request for Information to the county governments of Texas announcing funding to reduce hospitalizations and costs for adult PPH conditions by implementing interventions through a community coordinated approach. The 92 counties eligible to respond were those that had a hospitalization rate more than 50% higher than the state rate for at least one PPH condition from 2005-2009; and had a population of less than 100,000 residents between the ages of 18 and 64. DSHS was able to provide funding to 16 of the applicants: Angelina, Brooks, Ector, Grayson, Hunt, Liberty, Limestone, Nacogdoches, Orange, Polk, Red River, San Augustine, Tom Green, Trinity, Victoria, and Walker. The first contract period was January 1, 2012 – August 31, 2013.

DSHS re-contracted with 13 of the 16 counties for the period of September 1, 2013 – August 31, 2015. (Hunt, Liberty and Nacogdoches counties dropped out of the project.) With \$2 million available in Fiscal Years 2013 and 2014, the reduced number of funded counties added a condition of focus. The chart below shows conditions targeted by each of the 13 counties by August 2015:

Counties Contracting with DSHS	PPH Conditions County Projects Targeted prior to 9/1/15
Angelina	Bacterial Pneumonia, Dehydration, Hypertension, UTI
Brooks	Bacterial Pneumonia, Dehydration
Ector	Asthma, COPD, Diabetes
Grayson	Bacterial Pneumonia, COPD, Dehydration, UTI,
Limestone	Bacterial Pneumonia, CHF, COPD
Orange	Angina, Bacterial Pneumonia, CHF, COPD
Polk	Bacterial Pneumonia, CHF, COPD, Dehydration
Red River	CHF, COPD, Diabetes
San Augustine	Bacterial Pneumonia, Hypertension
Tom Green	Bacterial Pneumonia, COPD, Dehydration, UTI
Trinity	Bacterial Pneumonia, COPD, Hypertension, UTI
Victoria	Angina, Bacterial Pneumonia, CHF, Diabetes
Walker	Asthma, Hypertension, Diabetes

The PPH Initiative transferred from the Center for Policy and External Affairs to the Health Promotion and Chronic Disease Prevention Section in September 2015. The PPH Program operates with a funding level of \$2 million available for the 24-month period of September 1, 2015- August 31, 2017. The previously funded 13 counties continue participating. To make the greatest impact with limited resources, all counties are currently focusing on the highest-cost PPH conditions in Texas: CHF, COPD, and short and long-term diabetes complications. Each county is implementing four interventions to address the conditions: patient education, patient case management, healthcare provider education, and community education. Cross-cutting activities include immunizations, medication access, smoking

cessation, nutrition, physical activity, weight reduction, glycemic control, and blood pressure control.

The PPH Program is dedicated to providing technical assistance and support to local governments and other entities seeking to improve health care quality, reduce health care costs, and lower preventable hospitalization rates in order to improve the quality of life, health and wellbeing for Texans and their families. The PPH Surveillance Report is one of the tools available to advance our mission.

Definitions

1. **Bacterial Pneumonia** is an infection of the lungs that can cause mild to severe illness. It can often be prevented with vaccines and can usually be treated with antibiotics or specific drug therapies.
2. **Dehydration** means the body does not have enough fluid to function normally. Dehydration impacts older adults or institutionalized individuals who have a limited ability to communicate thirst. The condition commonly results from diarrhea and vomiting due to illness. People working outdoors in extreme heat conditions are also susceptible. Vulnerable people need to drink extra non-caffeinated fluids to keep from getting dehydrated. Mild fluid loss can most often be treated at home, but severe dehydration must be treated in the hospital.
3. **Urinary Tract Infection (UTI)** is usually caused when bacteria enter the bladder and cause inflammation and infection. UTIs are among the most common infections in people, and is usually treated with antibiotics. UTIs are the most common type of healthcare-associated infection. Among UTIs acquired in the hospital, approximately 75% are associated with a urinary catheter.
4. **Angina** (without procedures) is a symptom of coronary artery disease. Pain or discomfort in the chest, shoulders, arms, neck, jaw, or back; or a feeling like indigestion occurs because the heart muscle is not getting enough blood. Angina and other heart diseases can be prevented and treated by healthy lifestyle improvements and managing health conditions.
5. **Congestive Heart Failure (CHF)** happens when the heart cannot pump enough blood and oxygen to support other organs in the body. Early diagnosis and treatment can improve quality and length of life for people who have heart failure. Treatment usually involves **taking medications, reducing sodium** in the diet, and getting **daily physical activity**.
6. **Hypertension** (High Blood Pressure) is measured by the force of blood against your artery walls as it circulates through your body. Blood pressure normally rises and falls throughout the day, but it can cause health problems if it stays high for a long time such as heart disease and stroke. Hypertension can be controlled with prescribed medications and lifestyle changes.
7. **Chronic Obstructive Pulmonary Disease (COPD) or Asthma in Older Adults (age 40+)** refers to a group of diseases that cause airflow blockage and breathing-related problems. It includes emphysema, chronic bronchitis, and some cases of asthma. COPD treatment requires an individualized plan with multiple components that may include smoking cessation, medication, pulmonary rehabilitation, vaccination, oxygen supplements, etc.
8. **Diabetes** is a disease in which blood glucose levels are above normal. The pancreas makes a hormone called insulin to help glucose get into the body's cells. With diabetes, the body either doesn't make enough insulin or can't use its own insulin as well as it should, causing sugar to build up in the blood. A healthy food intake balanced by daily physical activities is the basic therapy for diabetes. Blood glucose levels must be closely monitored through frequent blood glucose testing. Insulin injections and/or oral medication are needed as well.

- a. **Short-term Complications** occur as a result of uncontrolled blood sugar levels. Ketoacidosis and hyperosmolarity occur from excessively high blood sugar levels. Dangerously high blood sugar (hyperglycemia) or dangerously low blood sugar (hypoglycemia) can lead to a coma.
- b. **Long-term Complications** are when renal, eye, neurological, circulatory, or not otherwise specified complications occur due to poor control of blood sugar levels over a period of time.

Note: definition sources are the Centers for Disease Control website and the Agency for Healthcare Research and Quality website. See **Related Links** on page 18.

Key Findings

- Females were disproportionately affected by hospitalizations for urinary tract infections. In Trinity County, females make up 51.7% of the population but accounted for 63.6% of potentially preventable hospitalizations for urinary tract infections.
- COPD or asthma in older adults had the highest risk-adjusted hospitalization rate compared to each of other reportable PPH conditions in Trinity County.
- COPD or asthma in older adults had the highest charge per adult Trinity County resident at \$284.
- Medicare was the most common primary payer source for each potentially preventable hospitalization condition with reportable data.

Demographics (Race/Ethnicity and Gender), Trinity County and Potentially Preventable Hospitalizations (PPH), 2014

Table 1 – Trinity County Demographics by Race/Ethnicity and Gender, 2014

	White Non-Hispanic		Black Non-Hispanic		Hispanic		Other		Total	
	N	%	N	%	N	%	N	%	N	%
Trinity	9,523	82.8	1,064	9.2	745	6.5	174	1.5	11,506	100.0

	Male		Female		Total	
	N	%	N	%	N	%
Trinity	5,556	48.3	5,950	51.7	11,506	100.0

Table 2 – PPH Demographics by Race/Ethnicity and Gender, 2014

PPH Conditions	White Non-Hispanic		Black Non-Hispanic		Hispanic		Other		Male		Female	
	N	%	N	%	N	%	N	%	N	%	N	%
Diabetes Short-term	--	--	--	--	--	--	--	--	--	--	--	--
Diabetes Long-term	--	--	--	--	--	--	--	--	--	--	--	--
COPD/Asthma in Older Adults	61	76.3	--	--	--	--	14	17.5	44	54.3	37	45.7
Hypertension	--	--	--	--	--	--	--	--	--	--	--	--
CHF	55	65.5	18	21.4	--	--	--	--	41	48.8	43	51.2
Dehydration	13	72.2	--	--	--	--	--	--	--	--	13	72.2
Bacterial Pneumonia	29	59.2	--	--	--	--	13	26.5	25	50.0	25	50.0
UTI	18	58.1	--	--	--	--	--	--	12	36.4	21	63.6
Angina	--	--	--	--	--	--	--	--	--	--	--	--

Note: The four highlighted conditions were selected for FY 2016 - 2017.

-- indicates less than 12 hospitalizations.

Please refer to the limitation section regarding reporting of race/ethnicity.

Figure 1 - Percentage of Trinity County Adults by Race/Ethnicity, 2014





Figure 2 - Percentage of PPH by Race/Ethnicity, Trinity County, 2014



Note: Data not reported for conditions with fewer than 12 hospitalizations reported for a specific race/ethnicity category. Please refer to the limitation section regarding reporting of race/ethnicity.

Demographics (Age Groups), Trinity County and Potentially Preventable Hospitalizations (PPH), 2014

Table 3 – Trinity County Demographics by Age Groups, 2014

	18-44 Years		45-64 Years		65-74 Years		75+ Years		Total	
	N	%	N	%	N	%	N	%	N	%
Trinity	3,622	31.5	4,334	37.7	2,116	18.4	1,434	12.5	11,506	100.0

Table 4 – PPH Demographics by Age Groups, Trinity County, 2014

PPH Conditions	18-44 years		45-64 years		65-74 years		75+ years	
	N	%	N	%	N	%	N	%
Diabetes Short-term	--	--	--	--	--	--	--	--
Diabetes Long-term	--	--	--	--	--	--	--	--
COPD/Asthma in Older Adults	--	--	27	33.3	22	27.2	31	38.3
Hypertension	--	--	--	--	--	--	--	--
CHF	--	--	12	14.3	24	28.6	43	51.2
Dehydration	--	--	--	--	--	--	--	--
Bacterial Pneumonia	--	--	13	26.0	--	--	29	58
UTI	--	--	--	--	--	--	15	45.5
Angina	--	--	--	--	--	--	--	--

Note: The four highlighted conditions were selected for FY 2016 - 2017.

-- indicates less than 12 hospitalizations.

COPD/Asthma in Older Adults is only among adults age 40 and over.

Number of Potentially Preventable Hospitalizations (PPH) and Risk Adjusted Rates, 2014

Definition of Risk Adjusted Rate: The risk adjusted rate allows us to compare communities that are different in their age and gender distribution. Risk adjusted rates statistically compensate (or adjust) for risk factor differences in two communities so that the outcome rates (here PPHs) can be compared legitimately despite the differences. In this report we have adjusted the rate for age and gender as those are identified as the risk factors that influence the PPHs.

For example, County A has 100 COPD hospitalizations, while County B has 200 COPD hospitalizations in the same year. This may lead to misinterpretation that County B has higher COPD hospitalizations. However, County A has a senior citizen population of 10% and County B has a senior citizen population of 40%. Using the risk adjusted rate we can compare these counties while accounting for the age differences in the population.


Table 5 - Number of PPH and the Risk Adjusted Rates, Trinity County, 2014

PPH Conditions	Number of Hospitalizations	Risk Adjusted Rates (95% Confidence Interval)
Diabetes Short-term	--	--
Diabetes Long-term	17	116.5 (63.5-169.4)
COPD/Asthma in Older Adults	81	800.2 (660.5-939.8)
Hypertension	15	105.1 (66.5-143.7)
CHF	84	520.7 (432.4-608.9)
Dehydration	18	117.0 (57.6-176.4)
Bacterial Pneumonia	50	319.4 (238.7-400.1)
UTI	33	225.1 (157.9-292.4)
Angina	--	--

Note: The four highlighted conditions were selected for FY 2016 - 2017.


-- indicates less than 12 hospitalizations.

Figure 3 – Risk Adjusted Rates of PPH, Trinity, 2014




Note: Data not reported for conditions with fewer than 12 hospitalizations reported.

Figure 4 – Risk Adjusted Rates of PPH for 4 Selected Conditions, Trinity County, 2014




Risk Adjusted Chronic Obstructive Pulmonary Disease (COPD) or Asthma in Older Adults Admission Rate Per 100,000 Adults by County, Texas, 2014



The risk adjusted COPD or asthma in older adults admission rate in Texas was 397.3 per 100,000 adults.

Data Classification: Quantiles


Risk Adjusted Diabetes Long-Term Complications Admission Rate Per 100,000 Adults by County, Texas, 2014



**The risk adjusted Diabetes Long-Term Complications admission rate
in Texas was 125.6 per 100,000 adults.**

Data Classification: Quantiles

Risk Adjusted Diabetes Short-Term Complications Admission Rate Per 100,000 Adults by County, Texas, 2014



**The risk adjusted Diabetes Short-Term Complications admission rate
in Texas was 66.2 per 100,000 adults.**

Data Classification: Quantiles

Hospital Charges and Length of Stay of Potentially Preventable Hospitalizations (PPH), Trinity County, 2014

Table 6 – Hospital Charges and Length of Stay of PPH, Trinity County, 2014

PPH Conditions	Average Charge (\$)	Median Charge (\$)	Total Charge (\$)	Charge per Adult County Resident (\$)	Average Length of Hospital Stay (Days)
Diabetes Short-term	--	--	--	--	--
Diabetes Long-term	48,020	28,278	816,348	71	6.2
COPD/Asthma in Older Adults	40,310	28,896	3,265,107	284	4.1
Hypertension	28,727	31,136	430,905	37	2.3
CHF	35,962	28,866	3,020,838	263	4.3
Dehydration	23,633	19,903	425,388	37	3.7
Bacterial Pneumonia	43,920	32,013	2,196,019	191	4.7
UTI	31,569	26,335	1,041,766	91	3.8
Angina	--	--	--	--	--

Note: The four highlighted conditions were selected for FY 2016 - 2017.

-- indicates less than 12 hospitalizations.

Expected Primary Source of Payment of Potentially Preventable Hospitalizations (PPH), Trinity County, 2014

Table 7 – Expected Primary Payer of PPH, Trinity County, 2014

PPH Conditions	Medicaid		Medicare		Private Health Insurance		Uninsured		Other	
	N	%	N	%	N	%	N	%	N	%
Diabetes Short-term	--	--	--	--	--	--	--	--	--	--
Diabetes Long-term	--	--	--	--	--	--	--	--	--	--
COPD/Asthma in Older Adults	--	--	62	76.5	--	--	--	--	--	--
Hypertension	--	--	--	--	--	--	--	--	--	--
CHF	--	--	72	85.7	--	--	--	--	--	--
Dehydration	--	--	--	--	--	--	--	--	--	--
Bacterial Pneumonia	--	--	37	74.0	--	--	--	--	--	--
UTI	--	--	25	75.8	--	--	--	--	--	--
Angina	--	--	--	--	--	--	--	--	--	--

Note: The four highlighted conditions were selected for FY 2016 - 2017.

-- indicates less than 12 hospitalizations.

Future Recommendations

- Incorporate the percentage of each PPH condition that has a secondary diagnosis of mental illness or substance abuse.
- Include more data visualizations, specifically the bar charts such as Figure 1 and 2 to compare between demographics with PPH conditions and county population demographics that further include age groups and gender.
- Develop benchmarks for condition-specific risk adjusted rates that allow for comparisons between the benchmark and county-level data.

Technical Notes

Measure Information:

The preventable hospitalization conditions in this report were selected from the Prevention Quality Indicators created by the Agency for Healthcare Research and Quality (AHRQ). AHRQ (www.ahrq.gov) is the lead federal agency responsible for research on health care quality, costs, outcomes, and patient safety.

Each potentially preventable hospitalization condition is defined by ICD-9-CM diagnosis codes. AHRQ may change the definition for these conditions each year.

All values may not add to the total because of missing data. The number of missing values may be different for each variable.

Limitations:

- Hospitalization data are based on inpatient hospitalization and do not include emergency department (ED) visits which did not result in hospital admission.
- The Texas Hospital Inpatient Discharge Public Use Data represent the number of inpatient hospitalizations. Since the data have been de-identified and an individual can be hospitalized more than once for the same condition during the data collection period, multiple hospitalizations for the same individual and the same diagnosis cannot be distinguished.
- Texas Hospital Inpatient Discharge Public Use Data is not a complete source of information on PPH-related discharges because some hospitals in Texas are exempt from reporting requirements. This may result in an underestimation of the number of related discharges.
- Incidence rates and patterns of newly diagnosed conditions are not captured by the data in this report. Using hospitalization data, we cannot determine if a hospitalization serves as an initial diagnosis of a specific condition.
- Hospital charges are based on all charges incurred during a hospital stay for a discharge where a specific PPH condition was the primary diagnosis. Charges may not be specifically associated with PPH care and may be associated with secondary diagnoses.
- Hospital charges are not the same as hospital costs or payments.
- Race and ethnicity data are generally not collected by hospitals and may be subjectively captured.
- Race is changed to 'Other' and ethnicity is suppressed if a hospital has fewer than ten discharges of a race in a single quarter.

Related Links:

- 1) PPH Program at Texas Department of State Health Services. (<http://www.dshs.state.tx.us/ph/>)
- 2) Prevention Quality Indicators, Agency for Healthcare Research and Quality. (http://www.qualityindicators.ahrq.gov/modules/pqi_resources.aspx)
- 3) Texas Hospital Inpatient Public Use Data File (PUDF), Texas Health Care Information Collection, Center for Health Statistics. (<http://www.dshs.state.tx.us/thcic/hospitals/inpatientpudf.shtm>)
- 4) Centers for Disease Control (CDC). (<http://www.cdc.gov/>)

Additional Information:

Health Promotion and Chronic Disease Prevention Section. (<https://www.dshs.state.tx.us/chronic/>)

